## DR ADRIAN SJARIF

MBBS (Hons), BSc, MS, FRACS

Provider No. 255103RF

ADRIAN SJARIF ABN 49 939 706 825 KOGARAH 14 KENSINGTON STREET KOGARAH NSW 2217 TELEPHONE: 9588 5700

MIRANDA SUITE 3, LEVEL 4 531 – 533 THE KINGSWAY MIRANDA NSW 2228 TELEPHONE: 9525 8400

WOLLONGONG 372 CROWN STREET WOLLONGONG NSW 2500 TELEPHONE: 4226 6111

AFTER HOURS: 0413 828 414

**Post Operative Instructions** 

## **FACELIFT**

- 1. Take all medications as directed
- 2. Sleep on 2 or 3 pillows to keep the head elevated. This will reduce swelling
- 3. Avoid bending down. Keep your head elevated at all times
- 4. Apply ice packs to the face as much as possible for the first 48 hours, e.g. for 20 minutes every 1-2 hours
- 5. REST ONLY. Do no exercise. Keep cool
- 6. Expect some swelling of the face. A small amount of blood ooze through the bandage can also be expected.

Further instructions will be given when we remove the dressings in two days. Please bring a scarf with you to this appointment

## **EYELID REDUCTION**

- 7. Use cold saline and cold water (boil first then chill) packs frequently to reduce swelling and soothe discomfort
- 8. Clean the eyes with a moist cotton bud 3 times a day
- 9. **If there is severe pain or visual impairment contact us immediately –** this would be an extremely unlikely occurrence.

For lubrication of the eyes (do not purchase until advised by surgeon):

Remove soft contact lenses before using

Day: "Refresh" or "Murine Revital Eyes" or similar eye lubricant

Night: "Viscotears Liquid Gel"

Follow up appointment:			
at:	☐ Kogarah Rooms	☐ Miranda Rooms	☐ Wollongong Rooms

If you have any questions or concerns, please contact us on one of the numbers above.

Please see the reverse for possible adverse reactions and drug interactions  $\rightarrow$ 

Possible adverse reactions and drug interactions:				
Medication	Possible Adverse Reaction			
Refresh Murine Revital Eyes	Stinging and burning			
/iscotears Liquid Gel	Mild transient burning, sticky eyelids, blurred vision			
	rse reaction/s to any of your medications, cease the medication and contact your Operative Instructions sheet) for further advice.			

